

Pulsar Timing in Extreme Mass Ratio Binaries

Tom Kimpson w/, Kinwah Wu, Silvia Zane.


July 10, 2019

Mullard Space Science Laboratory, UCL

A Problem

GR is incomplete

- Field equations = Non-unique
- Breaks down: Singularities + Quantum Gravity



- Extreme Mass Ratio Binary (EMRB)
- Event Horizon Clock

A precision, strong-field probe

$$\epsilon \propto \frac{M}{r}$$

Gravity Probe B



MICROSCOPE

Sun



Hulse-Taylor



Double PSR

S2



EMRB



Horizon

10^{-10}

10^{-8}

10^{-6}

10^{-4}

10^{-2}

10^0

ϵ

3 important parameters:

$$M, \chi, Q$$

Fundamental Physics

- No Hair Theorem ($Q = -\chi^2$)
- Cosmic Censorship Conjecture ($\chi \leq M^2$)

Astrophysics

- Astrophysical BH = Kerr solution?
- Constrain low end of $M - \sigma$ relation / Existence of IMBH

Hunting Grounds

- Galactic Centre
- Globular Clusters

Detection Prospects

- $\lesssim 10^4$ PSR at < 1 pc (Wharton et al. 2012, *ApJ* 753:108, Rajwade et al. 2017, *MNRAS* 471:730)
- Closest semi-major axis $\lesssim 100$ AU
- Closest pericentre distance ~ 2 AU (Zhang et al. 2014, *ApJ* 784:106)

- $\lesssim 10^4$ PSR at < 1 pc (Wharton et al. 2012, ApJ 753:108, Rajwade et al. 2017, MNRAS 471:730)
- Closest semi-major axis $\lesssim 100$ AU
- Closest pericentre distance ~ 2 AU (Zhang et al. 2014, ApJ 784:106)

No such PSR-EMRB yet detected!

Goal: Use the next-generation radio telescopes to time a pulsar in orbit around a massive central black hole.

Require theoretical basis for PSR **Timing Signal**

This Work: Why?

- Detection. *e.g. Are our algorithms good enough?*
- Modelling. *i.e. GR predictions vs. observation*

This Work: How?

Require theoretical basis for PSR **Timing Signal**


Behaviour of light + Orbital Dynamics = Timing signal

This Work: How?

Require theoretical basis for PSR **Timing Signal**

Behaviour of light + Orbital Dynamics = Timing signal

Ray Tracing



This Work: How?

Require theoretical basis for PSR **Timing Signal**

Behaviour of light + **Orbital Dynamics** = Timing signal

- Textbook GR: point particles.
- Real pulsars \neq point particles!

Creating the skeleton

$$T^{\mu\nu}{}_{;\nu} = 0$$

Multipole expansion to dipole order:

$$\frac{Dp^\mu}{d\tau} = -\frac{1}{2}R^\mu{}_{\nu\alpha\beta}u^\nu S^{\alpha\beta}$$

$$\frac{DS^{\mu\nu}}{d\tau} = p^\mu u^\nu - p^\nu u^\mu$$


(Mathisson 1937, Papetrou 1951, Dixon 1974)

Equations are
indeterminate


Spin Couplings

- Spin-spin
- Spin-orbit
- Spin-curvature



Orbital Dynamics: circular




Orbital Dynamics: eccentric



Spin Precession




Putting it all together...



Behaviour of light + Orbital Dynamics = Timing signal

Putting it all together...



Behaviour of light + Orbital Dynamics = Timing signal

Optimization problem

$$f(\alpha, \beta)$$

Effects to consider

- Gravitational lensing
- Primary/Secondary rays
- Influence of plasma: temporal/spatial dispersion
- Gravitational/Relativistic time dilation
- Orbital Dynamics
- Spin-curvature coupling (+spin-spin, spin-orbit)
- Spin precession
- Relativistic aberration


Photon ToA, pulse profile, intensity, observability

Effects to consider

- Gravitational lensing
- Primary/Secondary rays
- Influence of plasma: temporal/spatial dispersion
- Gravitational/Relativistic time dilation
- Orbital Dynamics
- Spin-curvature coupling (+spin-spin, spin-orbit)
- Spin precession
- Relativistic aberration


Photon ToA, pulse profile, intensity, observability

Gravitational Bending



- Deviation from straight lines
- Primary/Secondary Rays

Plasma: spatial dispersion



- PSR-EMRB = precision strong-gravity probes
- Require **fully relativistic $t - \nu$** model
- Open question: How good are current methods?

References:


Kimpson et al. 2019, doi:10.1093/mnras/stz138

Kimpson et al. 2019, doi:10.1093/mnras/stz845

Questions?

Putting it all together...

- Pulsar emission \neq isotropic
 - Find intersection with radiation point
- $$X_{\text{rad}}^i = R_{\text{PSR}} \hat{\mathbf{n}} + X_{\text{pulsar}}^i$$
- $\hat{\mathbf{n}} = \hat{\mathbf{n}}(S_\theta(\tau), \psi)$



Aberration

- 'Seen' if $\omega < \omega_c$
- Global $\omega \neq$ Local ω
- Transform to coming frame

